


Amendments To The Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

What is claimed is:

1. (Original) A compound of formula (I)



or a pharmaceutically acceptable salt, solvate, or derivative thereof, wherein:


X is a C₁₋₅ alkylene chain, wherein said X is optionally substituted by one or more =O, =S, -S(O)_t-, alkyl, or halogen and wherein said C₁₋₅ alkylene chain may optionally have 0-3 heteroatoms selected from oxygen, phosphorus, sulfur, or nitrogen;

Ring A is a saturated, partially saturated or aromatic 3-7 monocyclic or 8-10 membered bicyclic ring having one ring nitrogen and 0-4 additional heteroatoms selected from oxygen, phosphorus, sulfur, or nitrogen;


R¹ is selected from the group consisting of

(a) a saturated, partially saturated, or aromatic 4-7 monocyclic or 8-10 membered bicyclic ring having one ring nitrogen and 0-4 additional heteroatoms selected from oxygen, phosphorus, sulfur, or nitrogen, optionally attached through a C₁₋₆ alkylene chain, and optionally substituted by one or more R⁸;


(b)




(c)




(d)



(e)



(f)



Q is carbon, oxygen, or $-S(O)_t$;

w is 1 or 2;

each R^2 is independently selected from $-OR^0$, $-C(O)-R^0$, $-S(O)_2-R^0$, $-C(O)-N(R^0)_2$, $-S(O)_2-N(R^0)_2$, $-(CH_2)_a-N(R^0)(-V_b-R^+)$, $-(CH_2)_a-(-V_b-R^+)$, halogen, alkyl optionally substituted by one or more R^7 , alkenyl optionally substituted by one or more R^7 , alkynyl optionally substituted by one or more R^7 , aryl optionally substituted by one or more R^6 , heteroaryl optionally substituted by one or more R^6 , cycloalkyl optionally substituted by one or more R^8 , and heterocyclyl optionally substituted by one or more R^8 ; and two adjacent R^2 s on Ring A are optionally taken together to form a fused, saturated, partially saturated or aromatic 5-6 membered ring having 0-3 heteroatoms selected from oxygen, phosphorus, sulfur, or nitrogen; or two geminal R^2 s are optionally taken together to form a spiro, saturated, partially saturated or aromatic 5-6 membered ring having 0-3 heteroatoms selected from oxygen, phosphorus, sulfur, or nitrogen, said fused or spiro ring being optionally substituted by one or more R^8 ;

a is 0-3;

b is 0 or 1;

V is -C(O)-, -C(O)O-, -S(O)₂-, or -C(O)-N(R⁰)-;

R⁺ is alkyl, cycloalkyl, aralkyl, aryl, heteroaryl, heteroaralkyl, or heterocyclyl, wherein said R⁺ is optionally substituted by one or more R⁸;

d is 0-1;

m is 0 or 1;

n is 0-5;

each R³ independently is -H, -N(R⁰)₂, -N(R⁰)C(O)R⁰, -CN, halogen, -CF₃, alkyl optionally substituted by one or more groups selected from R⁷ or -S-aryl optionally substituted by -(CH₂)₁₋₆-N(R⁰)SO₂(R⁰), alkenyl optionally substituted by one or more groups selected from R⁷ or -S-aryl optionally substituted by -(CH₂)₁₋₆-N(R⁰)SO₂(R⁰), alkynyl optionally substituted by one or more groups selected from R⁷ or -S-aryl optionally substituted by -(CH₂)₁₋₆-N(R⁰)SO₂(R⁰), cycloalkyl or carbocyclyl optionally substituted by one or more R⁸, aryl optionally substituted by one or more R⁶, heteroaryl optionally substituted by one or more R⁶, or heterocyclyl optionally substituted by one or more R⁸;

Y is alkyl, alkenyl, alkynyl, -(CR⁴R⁵)_p-, -C(O)-, -C(O)C(O)-, -C(S)-, -O-(CH₂)₀₋₄-C(O)-, -(CH₂)₀₋₄-C(O)-O-, -N(R⁰)-C(O)-, -C(O)-N(R⁰)-, -N(R⁰)-C(S)-, -S(O)_t-, -O-C(=N-CN)-, -O-C(=N-R⁰)-, -C(=N-CN)-O-, -C(=N-CN)-S-, -C(=N-R⁰)-O-, -S-C(=N-CN)-, -N(R⁰)-C(=N-CN)-, -C(=N-CN)-, -N(R⁰)-C[=N-C(O)-R⁰]-, -N(R⁰)-C[=N-S(O)_t-R⁰]-, -N(R⁰)-C(=N-OR⁰)-, -N(R⁰)-C(=N-R⁰)-, or -C(=N-R⁰)-;

each R⁴ independently is H or alkyl optionally substituted by R⁷, alkenyl optionally substituted by R⁷, alkynyl optionally substituted by R⁷;

each R⁵ independently is selected from -H, -C(O)-OR⁶, -C(O)-N(R⁰)₂, -S(O)₂-N(R⁰)₂, -S(O)₂-R⁶, aryl optionally substituted by R⁶, or heteroaryl optionally substituted by R⁶;

p is 1-5;

each t independently is 1 or 2;

each R⁶ is independently selected from the group consisting of halogen, -CF₃, -OCF₃, -OR⁰, -(CH₂)₁₋₆-OR⁰, -SR⁰, -(CH₂)₁₋₆-SR⁰, -SCF₃, -R⁰, methylenedioxy, ethylenedioxy, -NO₂, -CN, -(CH₂)₁₋₆-CN, -N(R⁰)₂, -(CH₂)₁₋₆-N(R⁰)₂, -NR⁰C(O)R⁰, -NR⁰(CN), -NR⁰C(O)N(R⁰)₂, -NR⁰C(S)N(R⁰)₂, -NR⁰CO₂R⁰, -NR⁰NR⁰C(O)R⁰,

-NR⁰NR⁰C(O)N(R⁰)₂, -NR⁰NR⁰CO₂R⁰, -C(O)C(O)R⁰, -C(O)CH₂C(O)R⁰,
-(CH₂)₀₋₆CO₂R⁰, -O-C(O)R⁰, -C(O)R⁰, -C(O)N(R⁰)N(R⁰)₂, -C(O)N(R⁰)₂, -
C(O)N(R⁰)OH, -C(O)N(R⁰)SO₂R⁰, -OC(O)N(R⁰)₂, -S(O)_iR⁰, -S(O)_i-OR⁰,
-S(O)_iN(R⁰)C(O)R⁰,
-S(O)_iN(R⁰)OR⁰, -NR⁰SO₂N(R⁰)₂, -NR⁰SO₂R⁰, -C(=S)N(R⁰)₂, -C(=NH)-N(R⁰)₂,
-(CH₂)₁₋₆-C(O)R⁰, -C(=N-OR⁰)-N(R⁰)₂, -O-(CH₂)₀₋₆-SO₂N(R⁰)₂, -(CH₂)₁₋₆
NHC(O)R⁰, and -SO₂N(R⁰)₂ wherein the two R⁰s on the same nitrogen are
optionally taken together to form a 5-8 membered saturated, partially
saturated, or aromatic ring having additional 0-4 heteroatoms selected from
oxygen, phosphorus, nitrogen, or sulfur;

each R⁷ is independently selected from halogen, -CF₃, -R⁰, -OR⁰, -
OCF₃, -(CH₂)₁₋₆-OR⁰, -SR⁰, -SCF₃, -(CH₂)₁₋₆-SR⁰, aryl optionally substituted by -
R⁶, methylenedioxy, ethylenedioxy, -NO₂, -CN, -(CH₂)₁₋₆-CN, -N(R⁰)₂, -(CH₂)₁₋₆-
N(R⁰)₂, -NR⁰C(O)R⁰, -NR⁰(CN), -NR⁰C(O)N(R⁰)₂, -N(R⁰)C(S)N(R⁰)₂, -
NR⁰CO₂R⁰, -NR⁰NR⁰C(O)R⁰, -NR⁰NR⁰C(O)N(R⁰)₂, -NR⁰NR⁰CO₂R⁰,
-C(O)C(O)R⁰, -C(O)CH₂C(O)R⁰, -(CH₂)₀₋₆-CO₂R⁰, -C(O)R⁰, -C(O)N(R⁰)N(R⁰)₂, -
C(O)N(R⁰)₂, -C(O)N(R⁰)OH, -OC(O)R⁰, -C(O)N(R⁰)SO₂R⁰, -OC(O)N(R⁰)₂,
-S(O)_iR⁰, -S(O)_i-OR⁰, -S(O)_iN(R⁰)C(O)R⁰, -S(O)_iN(R⁰)OR⁰, -NR⁰SO₂N(R⁰)₂,
-NR⁰SO₂R⁰, -C(=S)N(R⁰)₂,
-C(=NH)-N(R⁰)₂, -(CH₂)₁₋₆-C(O)R⁰, -C(=N-OR⁰)-N(R⁰)₂, -O-(CH₂)₀₋₆-SO₂N(R⁰)₂,
-(CH₂)₁₋₆-NHC(O)R⁰, and -SO₂N(R⁰)₂ wherein the two R⁰s on the same nitrogen
are optionally taken together to form a 5-8 membered saturated, partially
saturated, or aromatic ring having additional 0-4 heteroatoms selected from
oxygen, phosphorus, nitrogen, or sulfur;

each R⁸ independently is selected from the group consisting of R⁷, =O,
=S, =N(R⁰), and =N(CN);

R⁹ is hydrogen, alkyl optionally substituted by one or more R⁷, alkenyl
optionally substituted by one or more R⁷, alkynyl optionally substituted by one
or more R⁷, cycloalkyl optionally substituted by one or more R⁸, heterocyclyl
optionally substituted by one or more R⁸, heteroaryl optionally substituted by
one or more R⁶, or aryl optionally substituted by one or more R⁶; or


-(Y)_m-R³ and R⁹ may combine with the nitrogen atom with which they are
attached to form a saturated, partially saturated, or aromatic 5-7 membered

monocyclic or 8-10 membered bicyclic ring that optionally contains 1 to 3 additional heteroatoms selected oxygen, phosphorus, sulfur, or nitrogen, wherein said ring may be optionally substituted with one or more R^8 ; R^{10} is hydrogen, alkyl optionally substituted by one or more R^7 , alkenyl optionally substituted by one or more R^7 , alkynyl optionally substituted by one or more R^7 , cycloalkyl optionally substituted by one or more R^8 , heterocyclyl optionally substituted by one or more R^8 , heteroaryl optionally substituted by one or more R^6 , or aryl optionally substituted by one or more R^6 ;


each R^0 is independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, carbocyclalkyl, aryl, heteroaryl, aralkyl, heteroaralkyl, heterocyclyl, and heterocyclalkyl, wherein each member of R^0 except H is optionally substituted by one or more R^* , $-OR^*$, $N(R^*)_2$, $=O$, $=S$, halogen, $-CF_3$, $-NO_2$, $-CN$, $-C(O)R^*$, $-CO_2R^*$, $-C(O)-aryl$, $-C(O)-heteroaryl$, aralkyl, $-S(O)_t-aryl$, $-S(O)_t-heteroaryl$, $-NR^*SO_2R^*$, $-NR^*C(O)R^*$, $-NR^*C(O)N(R^*)_2$, $-N(R^*)C(S)N(R^*)_2$, $-NR^*CO_2R^*$, $-NR^*NR^*C(O)R^*$, $-NR^*NR^*C(O)N(R^*)_2$, $-NR^*NR^*CO_2R^*$, $-C(O)C(O)R^*$, $-C(O)CH_2C(O)R^*$, $-C(O)N(R^*)N(R^*)_2$, $-C(O)N(R^*)_2$, $-C(O)NR^*SO_2R^*$, $-OC(O)N(R^*)_2$, $-S(O)_tR^*$, $-NR^*SO_2N(R^*)_2$, and $-SO_2N(R^*)_2$ wherein the two R^* s on the same nitrogen are optionally taken together to form a 5-8 membered saturated, partially saturated or aromatic ring having additional 0-4 heteroatoms selected from oxygen, phosphorus, nitrogen or sulfur; and

each R^* is independently H, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, or heteroaryl.


2. (Original) The compound of claim 1 wherein R^{10} is optionally substituted aryl.
3. (Original) The compound of claim 2 wherein R^{10} is optionally substituted phenyl.
4. (Original) The compound of claim 1 wherein R^1 is





5. (Original) The compound of claim 4 where R^9 is alkyl.
6. (Original) The compound of claim 5 wherein R^9 is methyl.
7. (Original) The compound of claim 4 wherein $-(Y)_m-R^3$ is selected from the group consisting of




8. (Original) The compound of claim 4 wherein $-(Y)_m-R^3$ is selected from the group consisting of











9. (Original) The compound of claim 4 wherein $-(Y)_mR^3$ and $-R^9$ combine with the nitrogen atom to which they are attached to form



10. (Currently Amended) The compound of claim 1 wherein R¹ is selected from




11. (Original) The compound of claim 1 wherein X is $-(CH_2)-$, $-(CH_2-CH_2)-$, or $-(CH_2-CH_2-CH_2)-$.


12. (Original) The compound of claim 9 wherein X is optionally substituted by one or more halogen or oxo.


13. (Original) The compound of claim 9 wherein X optionally has 1-3 heteroatoms selected from oxygen, phosphorus, sulfur, or nitrogen.


14. (Original) The compound of claim 1 wherein the A ring is selected from the group consisting of




15. (Original) The compound of claim 12 wherein each R², with an asterisk indicating a point of substitution from Ring A, independently is selected from:









16. (Original) The compound of claim 1 wherein ring A, with two geminal R^2 s, is selected from:







17. (Original) The compound of claim 1 wherein the A ring is tropane or piperidine, either optionally substituted with one or more R^2 .

18. (Original) The compound of claim 15 wherein the A ring in combination with R² is










19. (Original) The compound of claim 1 wherein the A ring contains at least one additional nitrogen atom.

20. (Original) The compound of claim 17 wherein said A ring optionally is N-substituted.

21. (Original) The compound of claim 18 wherein the A ring is N-substituted with $-(CH_2)_a-(V_b-R^+)$.

22. (Original) The compound of claim 1 wherein the compound of formula (I) is:



wherein X is a C₂-C₃ alkylene chain and R³ and R⁹ are each as defined in claim 1.

23. (Currently Amended) A method of treatment of a viral infection in a mammal comprising administering to said mammal an antiviral effective amount of a compound according to ~~claims 1-20~~ claim 1.

24. (Currently Amended) A method according to claim ~~24~~ 23 wherein the viral infection is an HIV infection.

25. (Currently Amended) A method of treatment of a bacterial infection in a mammal comprising administering to said mammal an effective amount of a compound according to ~~claims 1-20~~ claim 1.

26. (Currently Amended) A method according to claim ~~23~~ 25 wherein the bacterium is *Yersinia pestis*.

27. (Currently Amended) A method of treatment of multiple sclerosis, rheumatoid arthritis, autoimmune diabetes, chronic implant rejection, asthma, rheumatoid arthritis, Crohns Disease, inflammatory bowel disease, chronic inflammatory disease, glomerular disease, nephrotoxic serum nephritis, kidney disease, Alzheimer's Disease , autoimmune encephalomyelitis, arterial thrombosis, allergic rhinitis, arteriosclerosis, Sjogren's syndrome (~~dermatomyositis~~), systemic lupus erythematosus, graft rejection, cancers with leukocyte infiltration of the skin or organs, infectious disorders including bubonic and pneumonic plague, human papilloma virus infection, prostate cancer, wound healing, amyotrophic lateral sclerosis and immune mediated disorders in a mammal comprising administering to said mammal a pharmaceutically effective amount of a compound according to ~~claims 1-20~~ claim 1.

28. (Currently Amended) A compound according to ~~claims 1-20~~ claim 1 for use in medical therapy.

29. (Cancelled).

30. (Cancelled).

31. (Cancelled).

32. (Cancelled).

33. (Cancelled).

34. (Currently Amended) A pharmaceutical composition comprising a pharmaceutically effective amount of a compound according to ~~claims 1-20~~ claim 1 together with a pharmaceutically acceptable carrier.

35. (Currently Amended) The pharmaceutical composition according to claim ~~32~~ 34 in the form of a tablet or capsule.

36. (Currently Amended) The pharmaceutical composition according to claim ~~32~~ 34 in the form of a liquid.

37. (Currently Amended) A method of treatment of a viral infection in a mammal comprising administering to said mammal a composition comprising a compound according to ~~claims 1-20~~ claim 1 and another therapeutic agent.

38. (Currently Amended) The method according to claim ~~35~~ 37, wherein said composition comprises another therapeutic agent selected from the group consisting of (1-alpha, 2-beta, 3-alpha)-9-[2,3-bis(hydroxymethyl)cyclobutyl]guanine [(-)BHCG, SQ-34514, lobucavir], 9-[(2R,3R,4S)-3,4-bis(hydroxymethyl)-2-oxetanosyl]adenine (oxetanocin-G), acyclic nucleosides, acyclovir, valaciclovir, famciclovir, ganciclovir, penciclovir, acyclic nucleoside phosphonates, (S)-1-(3-hydroxy-2-phosphonylmethoxypropyl)cytosine (HPMPC), [[[2-(6-amino-9H-purin-9-yl)ethoxy]methyl]phosphinylidene] bis(oxymethylene)-2,2-dimethylpropanoic acid (bis-POM PMEA, adefovir dipivoxil), [(1R)-2-(6-amino-9H-purin-9-yl)-1-methylethoxy]methyl]phosphonic acid (tenofovir), (R)-[[2-(6-Amino-9H-purin-9-yl)-1-methylethoxy]methyl]phosphonic acid bis-(isopropoxycarbonyloxymethyl)ester (bis-POC-PMPA), ribonucleotide reductase inhibitors, 2-acetylpyridine 5-[(2-chloroanilino)thiocarbonyl]thiocarbonohydrazone and hydroxyurea, nucleoside reverse transcriptase inhibitors, 3'-azido-3'-deoxythymidine (AZT, zidovudine), 2',3'-dideoxycytidine (ddC, zalcitabine), 2',3'-dideoxyadenosine, 2',3'-dideoxyinosine (ddI, didanosine), 2',3'-didehydrothymidine (d4T, stavudine), (-)-beta-D-2,6-diaminopurine dioxolane (DAPD), 3'-azido-2',3'-dideoxythymidine-5'-H-

phosphonate (phosphonovir), 2'-deoxy-5-iodo-uridine (idoxuridine), (-)-cis-1-(2-hydroxymethyl)-1,3-oxathiolane 5-yl)-cytosine (lamivudine), cis-1-(2-(hydroxymethyl)-1,3-oxathiolan-5-yl)-5-fluorocytosine (FTC), 3'-deoxy-3'-fluorothymidine, 5-chloro-2',3'-dideoxy-3'-fluorouridine, (-)-cis-4-[2-amino-6-(cyclopropylamino)-9H-purin-9-yl]-2-cyclopentene-1-methanol (abacavir), 9-[4-hydroxy-2-(hydroxymethyl)but-1-yl]-guanine (H2G), ABT-606 (2HM-H2G) ribavirin, protease inhibitors, indinavir, ritonavir, nelfinavir, amprenavir, saquinavir, fosamprenavir, (R)-N-tert-butyl-3-[(2S,3S)-2-hydroxy-3-N-[(R)-2-N-(isoquinolin-5-yloxyacetyl)amino-3-methylthiopropionyl]amino-4-phenylbutanoyl]-5,5-dimethyl-1,3-thiazolidine-4-carboxamide (KNI-272), 4R-(4alpha,5alpha,6beta)]-1,3-bis[(3-aminophenyl)methyl]hexahydro-5,6-dihydroxy-4,7-bis(phenylmethyl)-2H-1,3-diazepin-2-one dimethanesulfonate (mozenavir), 3-[1-[3-[2-(5-trifluoromethylpyridinyl)-sulfonylamino]phenyl]propyl]-4-hydroxy-6alpha-phenethyl-6beta-propyl-5,6-dihydro-2-pyranone (tipranavir), N'-[2(S)-Hydroxy-3(S)-[N-(methoxycarbonyl)-l-tert-leucylamino]-4-phenylbutyl-N alpha-(methoxycarbonyl)-N'-[4-(2-pyridyl)benzyl]-L-tert-leucylhydrazide (BMS-232632), 3-(2(S)-Hydroxy-3(S)-(3-hydroxy-2-methylbenzamido)-4-phenylbutanoyl)-5,5-dimethyl-N-(2-methylbenzyl)thiazolidine-4(R)-carboxamide (AG-1776), N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenyl-methyl-4(S)-hydroxy-5-(1-(1-(4-benzo[b]furanylmethyl)-2(S)-N'-(tert-butylcarboxamido)piperazinyl)pentanamide (MK-944A), interferons, α -interferon, renal excretion inhibitors, probenecid, nucleoside transport inhibitors, dipyridamole, pentoxifylline, N-acetylcysteine (NAC), Procysteine, α -trichosanthin, phosphonoformic acid, immunomodulators, interleukin II, thymosin, granulocyte macrophage colony stimulating factors, erythropoetin, soluble CD₄ and genetically engineered derivatives thereof, non-nucleoside reverse transcriptase inhibitors (NNRTIs), nevirapine (BI-RG-587), alpha-((2-acetyl-5-methylphenyl)amino)-2,6-dichloro-benzeneacetamide (loviride), 1-[3-(isopropylamino)-2-pyridyl]-4-[5-(methanesulfonamido)-1H-indol-2-ylcarbonyl]piperazine monomethanesulfonate (delavirdine), (10R, 11S, 12S)-12-hydroxy-6,6,10,11-tetramethyl-4-propyl-11,12-dihydro-2H,6H,10H-benzo(1,2-b:3,4-b':5,6-b'')tripyrans-2-one ((+) calanolide A), (4S)-6-Chloro-4-[1E)-cyclopropylethenyl]-3,4-dihydro-4-(trifluoromethyl)-2(1H)-quinazolinone

(DPC-083), (S)-6-chloro-4-(cyclopropylethynyl)-1,4-dihydro-4-(trifluoromethyl)-2H-3,1-benzoxazin-2-one (efavirenz, DMP 266), 1-(ethoxymethyl)-5-(1-methylethyl)-6-(phenylmethyl)-2,4(1H,3H)-pyrimidinedione (MKC-442), and 5-(3,5-dichlorophenyl)thio-4-isopropyl-1-(4-pyridyl)methyl-1H-imidazol-2-ylmethyl carbamate (capravirine), glycoprotein 120 antagonists, PRO-2000, PRO-542, 1,4-bis[3-[(2, 4- dichlorophenyl)carbonylamino]-2-oxo-5,8-disodiumsulfanyl]naphthalyl-2, 5-dimethoxyphenyl-1, 4-dihydrazone (FP-21399), cytokine antagonists, reticulose (Product-R), 1,1'-azobis-formamide (ADA), 1,11-(1,4-phenylenebis(methylene))bis-1,4,8,11-tetraazacyclotetradecane octahydrochloride (AMD-3100), integrase inhibitors, and fusion inhibitors.

39. (Currently Amended) A method of treatment of a viral infection in a mammal comprising administering to said mammal a composition comprising a compound according to ~~claims 1-20~~ claim 1 and ritonavir.